

# Spin-resolved second-order correlation energy of the two-dimensional uniform electron gas

Michael Seidl, Institute of Theoretical Physics, University of Regensburg,  
D-93040 Regensburg, Germany

**Abstract.** For the two-dimensional electron gas, the exact high-density limit of the correlation energy is evaluated here numerically for all values of the spin polarization. The result is spin-resolved into  $\uparrow\uparrow$ ,  $\uparrow\downarrow$ , and  $\downarrow\downarrow$  contributions and parametrized analytically. Interaction-strength interpolation yields a simple model (LSD) for the correlation energy at finite densities.

In recent years, two-dimensional (2D) electron systems have become the subject of extensive research [1]. The 2D version of density functional theory (DFT) has proven particularly successful in studying quantum dots [2, 3, 4]. The local spin-density approximation (LSD) of DFT requires the correlation energy of the spin-polarized uniform electron gas. This quantity in 2D is known accurately for a wide range of densities and spin polarizations from fixed-node diffusion Monte Carlo simulations [5]. Its high-density limit is known exactly in terms of six-dimensional momentum-space integrals [6]. Resolved into contributions due to  $\uparrow\uparrow$ ,  $\uparrow\downarrow$ , and  $\downarrow\downarrow$  excitation electron pairs, these integrals are evaluated here numerically. The analytical parametrization of the results, Eqs. (16) and (17) below, is a crucial ingredient for the construction of the spin-resolved correlation energy at finite densities, performed recently for the 3D electron gas [7]. It is also required for studying the magnetic response of the spin-polarized 2D electron gas [8, 9]. Generally, it provides a fundamental test for numerical parametrizations of the correlation energy [5].

In the 2D uniform electron gas, the electrons are moving on a plane at uniform density  $\rho = [\pi(r_s a_B)^2]^{-1}$ , where  $a_B = 0.529 \text{ \AA}$  is the Bohr radius and  $r_s$  is the dimensionless density parameter (Seitz radius). We consider lowest-energy states with a given spin polarization

$$\zeta \equiv \frac{\rho_{\uparrow} - \rho_{\downarrow}}{\rho} \quad (1)$$

where  $\rho_{\uparrow}$  and  $\rho_{\downarrow} \equiv \rho - \rho_{\uparrow}$ , respectively, are the (uniform) densities of spin-up and spin-down electrons. Including a neutralizing positive background, the total energy per electron is a unique function of the dimensionless parameters  $r_s$  and  $\zeta$ ,

$$e_{tot}(r_s, \zeta) = t_s(r_s, \zeta) + e_x(r_s, \zeta) + e_c(r_s, \zeta). \quad (2)$$

The non-interacting kinetic and exchange energies,

$$t_s(r_s, \zeta) = \frac{1 + \zeta^2}{2} \frac{1}{r_s^2}, \quad e_x(r_s, \zeta) = -\frac{4\sqrt{2}}{3\pi} \frac{(1 + \zeta)^{3/2} + (1 - \zeta)^{3/2}}{2} \frac{1}{r_s} \quad (3)$$

(all energies are given in units of  $1 \text{ Ha} \equiv e^2/a_B = 27.21 \text{ eV}$  in the following), may be understood as the 0th- and the 1st-order terms of a perturbation expansion for the electron-electron interaction (where  $r_s$  turns out to be the expansion parameter).

The remaining *correlation energy* in Eq. (2) appears to have the perturbation (high-density) expansion [10, 11]

$$e_c(r_s, \zeta) = \sum_{n=0}^{\infty} [a_n(\zeta) \ln(r_s) + b_n(\zeta)] r_s^n \quad (r_s \ll 1). \quad (4)$$

For the 2D electron gas (but not for the 3D one), the first coefficient vanishes,  $a_0(\zeta) \equiv 0$ . Consequently, the second-order ( $n = 0$ ) term is  $e_c^{(2)}(\zeta) \equiv b_0(\zeta)$ , representing the high-density ( $r_s \rightarrow 0$ ) limit of  $e_c(r_s, \zeta)$ . It can be split into an exchange (“2b”) and a ring-diagram (“2r”) term [6],

$$e_c^{(2)}(\zeta) = e_c^{(2b)} + e_c^{(2r)}(\zeta). \quad (5)$$

The exchange term has only equal-spins contributions,  $e_c^{(2b)} = e_{c\uparrow\uparrow}^{(2b)}(\zeta) + e_{c\downarrow\downarrow}^{(2b)}(\zeta)$ , given by the  $\delta_{\sigma_1\sigma_2}$  term of Eq. (14) in Ref. [6] (we choose the  $k_x$  axis in the direction of  $\mathbf{q}$ ),

$$e_{c,\sigma\sigma}^{(2b)}(\zeta) = \frac{1}{8\pi^2} \int_0^\infty \frac{dq}{q} \int_{A[\kappa_\sigma(\zeta), q]} d^2k_1 \int_{A[\kappa_\sigma(\zeta), q]} d^2k_2 \frac{1}{|q\mathbf{e}_x + \mathbf{k}_1 + \mathbf{k}_2|} \frac{1}{q + k_{1x} + k_{2x}}. \quad (6)$$

Here,  $q$ ,  $\mathbf{k}_1$ , and  $\mathbf{k}_2$  are dimensionless,  $\sigma \in \{\uparrow, \downarrow\}$ , and the domain of the 2D integrals is

$$A[\kappa, q] \equiv \left\{ \mathbf{k} \in \mathbb{R}^2 \mid |\mathbf{k}| < \kappa, \quad |\mathbf{k} + q\mathbf{e}_x| > \kappa \right\}, \quad \kappa_\sigma(\zeta) \equiv [1 + \text{sgn}(\sigma)\zeta]^{1/2}. \quad (7)$$

[ $\kappa_\sigma(\zeta)$  is the Fermi wave vector for spin- $\sigma$  electrons in units of its value at  $\zeta = 0$ .] Scaling the integration variables by some constant  $\kappa$ ,  $q = \kappa Q$  and  $\mathbf{k} = \kappa \mathbf{K}$ , we have generally

$$\int_0^\infty \frac{dq}{q} \int_{A[\kappa, q]} d^2k f(q, \mathbf{k}) = \kappa^2 \int_0^\infty \frac{dQ}{Q} \int_{A[1, Q]} d^2K f(\kappa Q, \kappa \mathbf{K}). \quad (8)$$

Applying this rule to the integrals in Eq. (6), we find [6]

$$e_{c,\sigma\sigma}^{(2b)}(\zeta) = [1 + \text{sgn}(\sigma)\zeta] J^{(2b)}. \quad (9)$$

Consequently [6], the full second-order exchange term  $e_c^{(2b)} = e_{c\uparrow\uparrow}^{(2b)}(\zeta) + e_{c\downarrow\downarrow}^{(2b)}(\zeta) \equiv 2J^{(2b)}$  is  $\zeta$ -independent. A Monte Carlo integration yields

$$J^{(2b)} \equiv e_{c\uparrow\uparrow}^{(2b)}(0) = (57.15 \pm 0.05) \text{ mHa} \quad (1 \text{ mHa} = 10^{-3} \text{ Ha}). \quad (10)$$

The ring-diagram term  $e_c^{(2r)}(\zeta)$  is the remaining part of expression (14) in Ref. [6], with the contributions

$$e_{c,\sigma_1\sigma_2}^{(2r)}(\zeta) = -\frac{1}{8\pi^2} \int_0^\infty \frac{dq}{q^2} \int_{A[\kappa_{\sigma_1}(\zeta),q]} d^2k_1 \int_{A[\kappa_{\sigma_2}(\zeta),q]} d^2k_2 \frac{1}{q + k_{1x} + k_{2x}}. \quad (11)$$

The equal-spins terms ( $\sigma_1 = \sigma_2$ ) can be treated in the same way as the integral (6),

$$e_{c,\sigma\sigma}^{(2r)}(\zeta) = -[1 + \text{sgn}(\sigma)\zeta] J^{(2r)}, \quad J^{(2r)} = (76.69 \pm 0.03) \text{ mHa}. \quad (12)$$

The only non-trivial  $\zeta$ -dependence is in the opposite-spins term  $e_{c\uparrow\downarrow}^{(2r)}(\zeta) \equiv e_{c\downarrow\uparrow}^{(2r)}(\zeta)$ ,

$$e_{c\uparrow\downarrow}^{(2r)}(\zeta) = e_{c\uparrow\downarrow}^{(2r)}(0) [1 - f(\zeta)]. \quad (13)$$

By definition,  $f(0) = 0$ , and, since  $A[\kappa_\downarrow(1), q] = \emptyset$ ,  $f(1) = 1$ . Moreover,  $e_{c\uparrow\downarrow}^{(2r)}(0) = -J^{(2r)}$ . When the results of a Monte Carlo evaluation of  $f(\zeta)$  at different values of  $\zeta$  are compared with the functions  $f_\alpha(\zeta) \equiv [(1+\zeta)^\alpha + (1-\zeta)^\alpha - 2]/(2^\alpha - 2)$ , particularly good agreement (specially for  $\zeta \rightarrow 0$  and  $\zeta \rightarrow 1$ ) is found in the limit  $\alpha \rightarrow 1$  (Fig. 1a),

$$f(\zeta) = f_1(\zeta) + \delta f(\zeta), \quad f_1(\zeta) \equiv \frac{(1+\zeta) \ln(1+\zeta) + (1-\zeta) \ln(1-\zeta)}{2 \ln 2}. \quad (14)$$

[Note that  $f_\alpha(\zeta)$  also represents the  $\zeta$ -dependence of  $t_s$  ( $\alpha = 2$ ) and  $e_x$  ( $\alpha = \frac{3}{2}$ ) in Eq. (3).] The small deviation  $\delta f(\zeta)$  is accurately fitted by a polynomial (Fig. 1b)

$$\delta f(\zeta) \approx 0.0636 \zeta^2 - 0.1024 \zeta^4 + 0.0389 \zeta^6. \quad (15)$$

The small minimum of  $\delta f(\zeta)$  indicated by the numerical data (dots in Fig. 1b) at  $\zeta \approx 0.98$  is probably real, since a similar peculiarity is observed for the 3D electron gas (see the inset in Fig. 1 of Ref. [12]).

In summary, the second-order correlation energy  $e_c^{(2)}(\zeta) = e_c^{(2b)} + e_c^{(2r)}(\zeta)$  is

$$e_c^{(2)}(\zeta) \equiv e_{c\uparrow\uparrow}^{(2)}(\zeta) + 2e_{c\uparrow\downarrow}^{(2)}(\zeta) + e_{c\downarrow\downarrow}^{(2)}(\zeta) = [153.38 f(\zeta) - 192.46] \text{ mHa}, \quad (16)$$

where  $f(\zeta)$  is given by Eqs. (14) and (15). The spin resolution is fixed by

$$e_{c\uparrow\uparrow}^{(2)}(\zeta) \equiv e_{c\downarrow\downarrow}^{(2)}(-\zeta) = -(1 + \zeta) \times 19.54 \text{ mHa}. \quad (17)$$

$e_c^{(2)}(\zeta) \equiv e_c(0, \zeta)$  is the high-density limit of the general correlation energy  $e_c(r_s, \zeta)$ . To illustrate the relevance of this limit for finite densities ( $r_s > 0$ ), the present result can be used in the interaction-strength interpolation (ISI) of Ref. [13]. This approach does not require the higher-order ( $n \geq 1$ ) terms of the expansion (4) (which is expected

to have only a finite radius of convergence). Instead, information beyond the second order is taken from the low-density (strong-interaction or Wigner-crystal) limit of the exchange-correlation energy  $e_{xc} \equiv e_x + e_c$  (per electron),

$$e_{xc}(r_s, \zeta) \rightarrow \frac{a_\infty}{r_s} + \frac{b_\infty}{r_s^{3/2}} \quad (r_s \rightarrow \infty). \quad (18)$$

The coefficients [14]  $a_\infty \approx -1.1061$  and  $b_\infty \approx \frac{1}{2}$  are independent of  $\zeta$ , since any spatial overlap between two electrons is strongly suppressed in this limit, no matter whether their spins are parallel or not [15]. The resulting ISI expression for the exchange-correlation energy at finite densities reads [13]

$$e_{xc}^{ISI}(r_s, \zeta) = \frac{a_\infty}{r_s} + \frac{2X}{Y} \left[ (1+Y)^{1/2} - 1 - Z \ln \left( \frac{(1+Y)^{1/2} + Z}{1+Z} \right) \right]. \quad (19)$$

Using  $b_\infty = \frac{1}{2}$  and writing  $e_x(r_s, \zeta) = c_x(\zeta)/r_s$ , we have explicitly [13]

$$\begin{aligned} X(r_s, \zeta) &= \frac{-b_0(\zeta)}{[c_x(\zeta) - a_\infty]^2} \frac{1}{r_s}, \\ Y(r_s, \zeta) &= \frac{4b_0(\zeta)^2}{[c_x(\zeta) - a_\infty]^4} r_s, \\ Z(\zeta) &= \frac{-b_0(\zeta)}{[c_x(\zeta) - a_\infty]^3} - 1. \end{aligned} \quad (20)$$

Eq. (19) provides a simple explicit LSD,

$$E_{xc}^{LSD}[\rho_\uparrow, \rho_\downarrow] = \int d^2r \rho(\mathbf{r}) e_{xc}^{ISI}(r_s(\mathbf{r}), \zeta(\mathbf{r})), \quad (21)$$

for treating arbitrary 2D electron systems (also finite ones such as quantum dots) by the Kohn-Sham Equations of DFT. In Eq. (21),  $r_s(\mathbf{r}) = a_B^{-1}[\pi\rho(\mathbf{r})]^{-1/2}$  and  $\zeta(\mathbf{r}) = [\rho_\uparrow(\mathbf{r}) - \rho_\downarrow(\mathbf{r})]/\rho(\mathbf{r})$ .

In Fig. 2a, the ISI prediction  $e_c^{ISI}(r_s, \zeta) = e_{xc}^{ISI} - e_x$  for the correlation energy of the unpolarized uniform electron gas ( $\zeta = 0$ ) is compared with the accurate parametrization of the fixed-node diffusion Monte Carlo results in Ref. [5].  $e_c^{ISI}$  differs slightly from the latter by up to 4%. This mild deviation might be cured by including in the ISI a simple model for the next-order coefficient of expansion (4) [16]. In the high-density limit ( $r_s \rightarrow 0$ ), however, where the present result is exact, the parametrization in Ref. [5] has for  $0.7 < \zeta < 0.95$  a small positive deviation [5], shown in Fig. 2b.

## Figure captions:

Fig. 1. (a) Numerical results (dots) for the function  $f(\zeta)$  of Eq. (13) obtained by Monte Carlo integrations of expression (11) (with  $\sigma_1\sigma_2 = \uparrow\downarrow$ ) at selected values of  $\zeta$ . The analytical function  $f_1(\zeta)$  of Eq. (14) is plotted as a dashed curve. The solid curve represents the accurate fit  $f_1(\zeta) + \delta f(\zeta)$ , using Eq. (15) for  $\delta f(\zeta)$ . (b) The fit (15) (solid curve) compared to the true deviation (dots) of the Monte-Carlo-integration results from  $f_1(\zeta)$ .

Fig. 2. The correlation energy of Ref. [5] (dotted curves) versus the present ISI results (solid curves).

## References

- [1] E. Abrahams, S. V. Kravchenko, and M. P. Sarachik, Rev. Mod. Phys. **73**, 251 (2001).
- [2] S.M. Reimann and M. Manninen, Rev. Mod. Phys. **74**, 1283 (2002).
- [3] H. Jiang, H.U. Baranger, and W. Yang, Phys. Rev. B **68**, 165337 (2003).
- [4] H. Saarikoski, E. Räsänen, S. Siljamäki, A. Harju, M.J. Puska, R.M. Nieminen, Phys. Rev. B **67**, 205327 (2003).
- [5] C. Attaccalite, S. Moroni, P. Gori-Giorgi, and G. B. Bachelet, Phys. Rev. Lett. **88**, 256 601 (2002).
- [6] A.K. Rajagopal and J.C. Kimball, Phys. Rev. B **15**, 2819 (1977).
- [7] P. Gori-Giorgi and J.P. Perdew, Phys. Rev. B (to appear; cf. cond-mat/0305250).
- [8] J. Moreno and D.C. Marinescu, Phys. Rev. B **68**, 195210 (2003)
- [9] M. Polini and M.P. Tosi, Phys. Rev. B **63**, 045118 (2001).
- [10] W. Macke, Z. Naturforsch. A **5A**, 192 (1950).
- [11] W.J. Carr and A.A. Maradudin, Phys. Rev. **133**, A371 (1964).
- [12] G.G. Hoffman, Phys. Rev. B **45**, 8730 (1992).
- [13] M. Seidl, J.P. Perdew, and S. Kurth, Phys. Rev. Lett. **84**, 5070 (2000).
- [14] L. Bonsall and A.A. Maradudin, Phys. Rev. B **15**, 1959 (1977).
- [15] M. Seidl, J.P. Perdew, and S. Kurth, Phys. Rev. A **62**, 012502 (2000).
- [16] M. Seidl and J.P. Perdew (unpublished).




Figure 1: (a) Numerical results (dots) for the function  $f(\zeta)$  of Eq. (13) obtained by Monte Carlo integrations of expression (11) (with  $\sigma_1\sigma_2 = \uparrow\downarrow$ ) at selected values of  $\zeta$ . The analytical function  $f_1(\zeta)$  of Eq. (14) is plotted as a dashed curve. The solid curve represents the accurate fit  $f_1(\zeta) + \delta f(\zeta)$ , using Eq. (15) for  $\delta f(\zeta)$ . (b) The fit (15) (solid curve) compared to the deviation (dots) of the Monte-Carlo-integration results from  $f_1(\zeta)$ .




Figure 2: The correlation energy of Ref. [5] (dotted curves) versus the present ISI results (solid curves).